

Reparil® Gel

SCHEDULING STATUS

S1

PROPRIETARY NAME AND DOSAGE FORM

Reparil® Gel

COMPOSITION

100 g contains

Aescin 1,0 g

Diethylamine salicylate 5,0 g

PHARMACOLOGICAL CLASSIFICATION

A 13.9 Dermatological preparations - special combinations

PHARMACOLOGICAL ACTION

Reparil® Gel has analgesic properties.

INDICATIONS

Symptomatic relief of pain associated with superficial soft tissue injuries, such as certain sports injuries.

Provides temporary relief from the minor pain and stiffness of rheumatism and muscle soreness.

CONTRAINDICATIONS

Hypersensitivity to the ingredients. Do not apply to broken skin and excessively raw areas.

DOSAGE AND DIRECTIONS FOR USE

Unless otherwise directed, a thin layer of the gel should be applied to the skin over the affected area several times daily.

SIDE EFFECTS AND SPECIAL PRECAUTIONS

Erythema. Desquamation. Urticaria and angio-neurotic oedema may occur in aspirin sensitive patients following exposure to local application.

Due to its alcohol content, **Reparil® Gel** should not be applied to areas of eczema or ulceration, nor to skin exposed to therapeutic irradiation, but only to the surrounding parts in order to avoid a possible burning sensation.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT

Systemic poisoning may occur from application to very large areas of skin.

IDENTIFICATION

A transparent, colourless to light yellow gel with odour of lavender and isopropyl alcohol.

PRESENTATION

Tubes of 20 g, 40 g and 100 g.

STORAGE INSTRUCTIONS

Keep the tube tightly closed. Store at or below 25 °C.

KEEP OUT OF REACH OF CHILDREN

REGISTRATION NUMBER

G/13.9/2367

NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION

XIXIA PHARMACEUTICALS (PTY) LTD

4 Brewery Street, Isando, Kempton Park, 1600

Republic of South Africa

DATE OF PUBLICATION OF THE PACKAGE INSERT

23 April 1993

Namibia Reg. No.: 05/13.9/0398

Zimbabwe Reg. No.: 79/14.19/1256

 **Mylan**

www.mylansa.co.za

56ZA2241321-00

706941



Reparil® Gel

SKEDULERINGSSTATUS

S1

EIENDOMSNAAM EN DOSEERVORM

Reparil® Gel

SAMESTELLING

100 g bevat

Eskien..... 1,0 g

Diëtielamiensilisilaat 5,0 g

FARMAKOLOGIESE KLASSIFIKASIE

A 13.9 Velpreparate-Besondere samestellings

FARMAKOLOGIESE WERKING

Reparil® Gel het analgetiese eienskappe.

INDIKASIE

Simptomatiese verligting van pyn wat gepaard gaan met oppervlakkige sagteweefselbeserings, soos met sekere sportbeserings.

Verskaf tydelike verligting van die mindere pyn en styfheid van rumatiek en spierseerheid.

KONTRA-INDIKASIES

Hipersensitiwiteit vir die bestanddele. Moet nie op stukkende vel of baie rou areas aangewend word nie.

DOSIS AND GEBRUIKSAANWYSINGS

Tensy anders voorgeskryf, moet 'n dun lagie van die jel 'n paar keer per dag op die vel oor die aangetaste deel aangewend word.

NEWE-EFFEKTE EN SPESIALE VOORSORGMATREËLS

Eriteem. Afskilfering. Urtikaria en angio-neurotiese edeem kan voorkom by aspirien-gevoelige pasiënte, na blootstelling aan direkte aanwending. Omdat dit alkohol bevat

en moontlik 'n branderige gevoel kan veroorsaak, mag **Reparil® Gel** nie gebruik word op ekseemagtige of ulkusvormende dele, of op velareas wat aan terapeutiese bestraling blootgestel was nie, maar slegs op die dele daaromheen.

BEKENDE SIMPTOME VAN OORDOSERING EN BESONDERHEDE VAN DIE BEHANDELING DAARVAN

Sistemiese vergifting kan voorkom wanneer die jel op baie groot velareas aangewend word.

IDENTIFIKASIE

'n Deursigtige, kleurlose tot liggeel jel met 'n laventel en isopropielalkohol geur.

AANBIEDING

Buise met 20 g, 40 g en 100 g.

BERGINGSAAWYSINGS

Maak die buis dig toe. Bewaar by of onder 25 °C.

HOU BUIE BEREIK VAN KINDERS

REGISTRASIENOMMER

G/13.9/2367

NAAM EN BESIGHEIDSADRES VAN DIE HOUER VAN DIE REGISTRASIE SERTIFIKAAT

XIXIA PHARMACEUTICALS (EDMS) Bpk.

4 Brewery Straat, Isando, Kempton Park, 1600
Republiek van Suid-Afrika

DATUM VAN PUBLIKASIE VAN DIE VOUBILJET

23 April 1993

Namibië Reg. Nr.: 05/13.9/0398

Zimbabwe Reg. Nr.: 79/14.19/1256

 **Mylan**

www.mylansa.co.za

56ZA2241321-00

706941

